

# How to Prescribe

**phexx®**

(lactic acid, citric acid, and  
potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

Formerly Phexxi



PATIENT: \_\_\_\_\_

DATE: \_\_\_\_\_

## Phexx Vaginal Gel (5g applicator)

**Sig:** Dispense Phexx or Phexxi. Administer 1 applicator intravaginally immediately before or up to 1 hour before EACH act of vaginal intercourse as needed.

**Dispense:** 30 Day Supply

- 180g (36 applicators)
- 120g (24 applicators)
- 60g (12 applicators)

---

NDC: 69751-101-12

REFILL: 11

## IMPORTANT SAFETY INFORMATION

### WARNINGS AND PRECAUTIONS

Few cases (0.36%) of adverse reactions of cystitis, pyelonephritis and other upper urinary tract infection (UTI) have been reported in Phexx clinical studies. Of these, one case of pyelonephritis was considered serious and required hospitalization. Avoid use of Phexx in females with history of recurrent urinary tract infection or urinary tract abnormalities.

**Please see additional Important Safety Information on reverse side and scan QR code for full Prescribing Information for Phexx.**



Formerly Phexxi

(lactic acid, citric acid, and potassium bitartrate) Vaginal Gel  
1.8%, 1%, 0.4%

## IMPORTANT SAFETY INFORMATION (cont'd)

### ADVERSE REACTIONS

Most common adverse reactions were vulvovaginal burning sensation, pruritus, mycotic infection, genital discomfort (including male partners); urinary tract infection, bacterial vaginosis, vaginal discharge, and dysuria. Patients should consult their healthcare provider for severe or prolonged genital irritation or urinary tract symptoms.

### Advise the patient:

- **To discontinue Phexx if they develop a local hypersensitivity reaction.**
- **That Phexx does not protect against HIV infection or other sexually transmitted infections.**
- **Avoid Phexx use with vaginal rings.**

**To report SUSPECTED ADVERSE REACTIONS, contact Evofem at toll-free phone**

**1-833-EVFMBIO or you may contact FDA at 1-800-FDA-1088 or [www.fda.gov/medwatch](http://www.fda.gov/medwatch).**

### INDICATIONS AND USAGE

Phexx is indicated for the prevention of pregnancy in females of reproductive potential for use as an on-demand method of contraception.

### LIMITATIONS OF USE

Phexx is not effective for the prevention of pregnancy when administered after intercourse.

**Please see additional  
Important Safety Information on  
reverse side and scan QR code for  
full Prescribing Information.**

